arXiv:1504.04001v1 [math.RA] 15 Apr 2015

TRIANGULABLE LEIBNIZ ALGEBRAS

TIFFANY BURCH AND ERNIE STITZINGER

ABSTRACT: A converse to Lie's theorem for Leibniz algebras is found and generalized. The result is used to find cases in which the generalized property, called triangulable, is 2-recognizeable; that is, if all 2-generated subalgebras are triangulable, then the algebra is also. Triangulability joins solvability, supersolvability, strong solvability, and nilpotentcy as a 2-recognizeable property for classes of Leibniz algebras.

Simultaneous triangulation of a set of linear transformations related to Leibniz algebras will be considered in this work. A necessary condition is that the minimum polynomials of the linear transformations are the products of linear factors. If the matrices commute, then this condition is sufficient. Lie's theorem is a generalization of this result, a result that is field dependent even in the algebraically closed case. Lie's theorem extends to Leibniz algebras. It can also be useful to have a condition on the set which guarantees simultaneous triangulation when extending to the algebraic closure, K, of the field of scalars, F. We will call this property, borrowing from Bowman, Towers and Varea [3], triangulable. Thus a Leibniz algebra A over a field F is triangulable on module M if the induced representation of $K \bigotimes A$ on $K \bigotimes M$ admits a basis such that the representing matrices are upper triangular. We find a condition on the Leibniz algebra that is equivalent to being triangulable. The condition is independent of the field and includes Lie's theorem. The condition is on A although triangulable refers to the action in the algebra that is over K. As an application, we consider to what extent an algebra is triangulable if all 2-generated subalgebras have the property. Such a property is called 2-recognizable, a concept investigated in groups, Lie algebras and Leibniz algebras for classes such as solvable, supersolvable, strongly solvable and, trivially, nilpotent and abelian. For an introduction to Leibniz algebras, see [1], [6] and [7].

Let M be a module for the Leibniz algebra A. Let left and right multiplication by $x \in A$ be denoted by T_x and S_x, respectively. If T_x is nilpotent, then S_x is also nilpotent and an Engle theorem holds; that is, A acts nilpotently on M [2]. Define A to be nil on M if this property holds. If I is an ideal of a Leibniz algebra A, then I is nil on A if and only if I is nilpotent. We will show that A is triangulable on itself if and only if A^2 is nilpotent. When A^2 is nilpotent, then A is called strongly solvable, a term introduced in [3].

Lemma 1. *Let B be an irreducible submodule of the A-module M such that dim$(B) = 1$. Then $A^2 \subseteq C_A(B) = \{a \in A | aB = 0 = Ba\}$.*

Proof. Let $x, y \in A$, $b \in B$ with $xb = \alpha_x b$ where α_x is a scalar. Then $(xy)b = x(yb) - y(xb) = x(\alpha_y b) - y(\alpha_x b) = \alpha_x \alpha_y b - \alpha_y \alpha_x b = 0$. Since B is irreducible, it follows that $b(xy) = 0$ ([1]). Hence $A^2 B = BA^2 = 0$ and the result holds. ☐

Theorem 1. *Let A be a Leibniz algebra and M be an A-module. There is a basis for M such that the representing matrices are in upper triangular form if and only if the linear transformations in the representation have minimum polynomials with all linear factors and A^2 is nil on M.*

We then have

Corollary 1. *Let A be a Leibniz algebra over an algebraically closed field and M be an A-module. There is a basis for M such that the representing matrices are in upper triangular form if and only if A^2 is nil on M.*

Proof of Theorem. The condition on the minimum polynomials is equivalent to being able to triangulate all the linear transformations. Suppose also that A^2 is nil on M. Then T_x, and hence S_x, is nilpotent for all $x \in A^2$ ([2]) and A^2 acts nilpotently on M by Engel's theorem. Let B be an irreducible submodule of M. It is enough to show that $\dim(B) = 1$ for then the result will follow by induction. Since the submodule $BA^2 + A^2B$ is contained in B, $BA^2 + A^2B$ is 0 or B. If the latter holds, then $B = BA^2 + A^2B = (BA^2 + A^2B)A^2 + A^2(BA^2 + A^2B) = \dots$, which is never 0, a contradiction. Hence $BA^2 + A^2B = 0$ and $BA^2 = 0 = A^2B$. Therefore, $0 = (xy)b = x(yb) - y(xb)$ for all $x, y \in A$, $b \in B$. Hence $x(yb) = y(xb)$ and $T_xT_y = T_yT_x$. Since B is irreducible, either $S_x = 0$ for all $x \in A$ or $S_x = -T_x$ for all $x \in A$ [1]. Then either $T_xS_y = -T_xT_y = -T_yT_x = S_yT_x$ or $T_xS_y = 0 = S_yT_x$ on B and $S_xS_y = T_xT_y = T_yT_x = S_yS_x$ or $S_xS_y = 0 = S_yS_x$ on B. Hence all T_x, S_y commute on B. Thus these transformations are simultaneously triangularizeable on B. Hence $\dim(B) = 1$ and the result holds in this direction.

Suppose that there is a basis for M such that all the transformations are triangularizeable on this basis. Then there is a one dimensional submodule B such that $A^2B = 0 = BA^2$ by the lemma. A^2 is nil on M/B by induction and hence on M. Since each transformation is triangularizeable, the minimum polynomials have linear factors. ☐

We state these results when A is acting on itself. A is called strongly solvable if A^2 is nilpotent ([3], [5]). A is supersolvable if there is an increasing chain of ideal of A, each of codimension 1 in the next.

Theorem 2. *Let A be a Leibniz algebra over an algebraically closed field. Then A is supersolvable if and only if A is strongly solvable.*

Proof. A is supersolvable if and only if A triangulable on itself. By Corollary 1, this is equivalent to A^2 being nil on A which in turn is the same as A^2 being nilpotent. The last statement is the definition of strongly solvable. ☐

The general case looks like the following.

Theorem 3. *A is triangulable if and only if A^2 is nil on A.*

Proof. Let $A^* = K \bigotimes A$ where K is the algebraic closure of the base field for A. If A^2 is nil on A, then A^2 is nilpotent as is $(A^2)^* = (A^*)^2$. Hence A^* is supersolvable by the last result and A is triangulable by definition. The converse holds by reversing the steps. ☐

A property is n-recognizeable in a class of algebras if whenever all n generated subalgebras have the property, then the algebra has the property. The concept when $n = 2$ has been considered in groups, Lie algebras and Leibniz algebras for classes that are solvable, supersolvable, nilpotent and strongly solvable (see [4], [5], [8]). A version of this has also been considered for triangulable in the Lie case in [3]. We considered it as an application of our result.

Theorem 4. *In the class of solvable Leibniz algebras, triangulable is 2 recognizable.*

Proof. Let N be a 2-generated subalgebra of A. Since N is triangulable, N^2 is nil on N and, thus, N is strongly solvable. Therefore A is strongly solvable by [5] and A^2 is nil on A. Then, using Theorem 3, A is triangulable. □

The result can be extended from solvable to all algebras in characteristic 0, again using a result in [5].

Corollary 2. *For Leibniz algebras, triangulable is 2-recognizeable at characteristic 0.*

Proof. Let A be a minimal counterexample. Simple Leibniz algebras are Lie algebras. Simple Lie algebras are 2-generated, hence A is triangulable. If I is a proper ideal in A, then all 2-generated subalgebras of I and A/I are strongly solvable by Theorem 3 and I and A/I are strongly solvable by Theorem 3 of [5] and A is solvable. The result holds by Theorem 4. □

References

[1] D. Barnes, Some theorems on Leibniz algebras, Comm. Alg. 39 (2011) 2463-2472

[2] L. Bosko, A. Hedges, J.T. Hird, N. Schwartz, K. Stagg, Jacobson's refinement of Engel's theorem for Leibniz algebras, Involve 4 (2011)293-296

[3] K. Bowman, D. Towers, V. Varea, Two generator subalgebras of Lie algebras, Linear and Multilinear Algebra 55 (2007) 429-438

[4] R. Brandl, Zur Theorie der untergruppenabgeschlossenen Formationen: endliche Varietten, J.Algebra 73 (1981) 1-22

[5] T. Burch, M. Harris, A. McAlister, E. Rogers, E. Stitzinger, S.M. Sullivan, 2-recognizeable classes of Leibniz algebras, J. Algebra. 423 (2015). 506-513

[6] I. Demir, K. Misra, E. Stitzinger, On some structures of Leibniz algebras, AMS Contemporary Math. 623 (2014) 41-54

[7] J.-L. Loday, Une version non commutative des algebres de Lie: les algebres de Leibniz, Enseign Math. (2) (1993) 39 (3-4) 269-293

[8] K. Moneyhun, E. Stitzinger, Some finite varieties of Lie algebras, J. Algebra 143 (1991) 173-17